

Gabrielle Whiting Letter to The Australian Cancer Council.

Gabrielle Whiting's letters are superbly well-written and extensively researched. There is just one correction. The U.S. National Cancer Institute has not changed its position that the abortion-breast cancer link is "non-existent." In the study, Dolle et al. 2009, National Cancer Institute branch chief, Louise Brinton, and colleagues at the Fred Hutchinson Cancer Research Center admitted that "...induced abortion and oral contraceptive use were associated with an increased risk for breast cancer."

29 January 2010

Professor Ian Olver  
Member Advisory Council  
Cancer Australia  
&  
Chief Executive Officer  
Cancer Council of Australia  
GPO Box 4708  
Sydney NSW 2001

Dear Professor Olver,

**N.C.I. (U.S.) Admission of Induced Abortion as a Risk Factor for Breast Cancer**  
**Urgent Action Required by Australian Cancer Authorities**

Regardless of -

- (a) an abundance of published peer-reviewed epidemiological evidence, that has accumulated since 1957, for an increased risk of breast cancer with induced abortion (including over 18 studies finding a statistically significant increase in risk<sup>1-18</sup>),
- (b) the findings of an actuary using mathematical formulae that, of seven known risk factors, induced abortion is the best predictor of breast cancer (the author of the study used the same mathematical model for a previous forecast that proved accurate for predicting cancers observed in earlier years),<sup>19</sup>
- (c) the recent publication of case control studies originating outside the purview of the National Cancer Institute (U.S.) (N.C.I.) on groups in Turkey and China, finding a statistically significant increase in risk of breast cancer with induced abortion,<sup>20,21</sup>
- (d) the scientifically valid refutation of flawed studies that claimed to repudiate the evidence for the induced abortion / breast cancer risk<sup>22-24</sup> (see also Professor Joel Brind's analysis of the Beral et al collaborative [Oxford] study [*The Lancet* 2004] on the Breast Cancer Prevention Institute's web site<sup>25</sup>),
- (e) the published analyses by various experts of the rejection of the valid science finding increased risk of breast cancer with induced abortion,<sup>26-28</sup>
- (f) the exposure of a fraudulent 2008 study, co-funded by the N.C.I. and co-authored by a member of the N.C.I., that claimed to refute the link,<sup>29-31</sup>
- (g) a plausible and unrefuted biological explanation for the link,<sup>32,33</sup>
- (h) the successful suing of abortion providers in U.S. and Australia for failure to inform women of the risk (in which the defendants negotiated out-of-court settlements to prevent the scientific evidence of the abortion / breast cancer risk being aired in Court),
- (i) the success of information campaigns by Endeavour Forum Inc. and others<sup>a</sup>, including web sites such as that of the Breast Cancer Prevention Institute<sup>34</sup> and YouTube videos<sup>35</sup> (that have reached many Australians including politicians, the public and employees of print and broadcast media), on the evidence for an increased risk of breast cancer with induced abortion

- the N.C.I. has hitherto denied that induced abortion increases the risk of breast cancer.

Australian cancer authorities have failed to acknowledge the strength of the evidence and ensure that women are informed of this serious risk for a terminal disease, typically advising to the effect that these studies must be treated with caution! Caution towards whom? The hapless women who may be exposed to a terminal, mutilating disease?

In a recent published peer-reviewed study finding an association between oral contraceptive use and triple negative breast cancer in women under 45, that is co-authored by Dr Louise Brinton of the N.C.I. who had previously influenced the N.C.I. to deny the abortion / breast cancer risk, **the N.C.I. has now admitted increased risk of breast cancer with induced abortion.**<sup>36</sup> The study's results included "Specifically, older age, family history of breast cancer, earlier menarche age, induced abortion, and oral contraceptive use were associated with an increased risk for breast cancer."

Table 1 in the published study listed induced abortion under "known and suspected risk factors" and revealed a statistically significant 40% risk increase for women who have had induced abortions.

Australian cancer authorities must avoid creating a perception in the community, which has become increasingly well-informed of the issue, that they are not dealing with the induced abortion / breast cancer risk because of a desire to preserve the status of abortion as a safe procedure.

Endeavour Forum Inc. believes we act on behalf of all Australian women when we call for Cancer Australia, the Cancer Council of Australia and the Cancer Council Victoria to revise their previous position on this issue and, without further delay, inform the public and conduct a dedicated education campaign on this serious risk for a terminal disease.

As a corollary to this we recommend that abortionists be required (a) to include advice in abortion consent forms that women increase their risk of breast cancer with induced abortion and (b) to personally counsel women requesting abortion on the risk.

Women will be assisted much more by being empowered to make informed choices with respect to breast cancer risks than by all the daffodil days, pink ribbon days and jumping around in pink suits.

The failure to date to inform women of the increased risk of breast cancer with induced abortion is scandalous. Any further delay in informing the public and the Australian and State and Territory Governments of this risk would be a scandal of even greater enormity. Thousands more Australian women may die or suffer treatment for the early progress of the disease and mutilation. The evidence has increased, as have admissions by medical and cancer authorities including the key N.C.I.<sup>b</sup>, to a point at which it is simply impossible for it to be withheld from the Australian public.

Yours sincerely,

Gabrielle Whiting  
Research Officer  
Endeavour Forum Inc.

cc: Professor David Hill, Director, Cancer Council Victoria  
The Hon Nicola Roxon MP, Minister for Health and Ageing, Australian Government  
The Hon Peter Dutton MP, Shadow Minister for Health and Ageing

<sup>a</sup>See copy enclosed of "In denial over the abortion/breast cancer link" by Terri Kelleher, *Journal of the Australian Family Association* 2009;30(3):30-37.

<sup>b</sup>Cancer institutes and medical groups that recognize the link include:-

Breast Cancer Prevention Institute, New Jersey, U.S.A; National Physicians Center for Family Resources, Birmingham, Alabama, U.S.A; Catholic Medical Association U.S.A.; American Association of Pro-Life Obstetricians and Gynecologists; The Polycarp Research Institute, Altoona, Pennsylvania, U.S.A.; Ethics and Medics, Philadelphia, Pennsylvania, U.S.A.; MaterCare International, Newfoundland, Canada; Breast Care Center-EAMC, Quezon City, Philippines. The Association of American Physicians and Surgeons supports disclosure of research supporting the induced abortion / breast cancer link.

## References

1. Segi M, Fukushima I, Fujisaku S et al. An epidemiological study on cancer in Japan. *GANN (Jpn. J. Cancer Res.)* 48 suppl. 1957:1-63.
2. Rosenberg L, Palmer JR, Kaufman DW et al. Breast cancer in relation to the occurrence and time of induced and spontaneous abortion. *Am. J. Epidemiol.* 1988 May;127(5):981-9.
3. Howe HL, Senie RT, Bzduch H et al. Early abortion and breast cancer risk among women under age 40. *Int. J. Epidemiol.* 1989;18:300-304.
4. Laing AE, Demenais FM, Williams R et al. Breast cancer risk factors in African-American women: The Howard University Tumor Registry experience. *J. Natl. Med. Assoc.* 1993;85(12):931-9.
5. Laing AE, Bonney GE, Adams-Campbell L et al. Reproductive and lifestyle factors for breast cancer in African-American women. *Genet. Epidemiol.* 1994;11:300 (abstract).
6. Daling JR, Malone DE, Voigt LF, White E, Weiss NS. Risk of breast cancer among young women: Relationship to induced abortion. *J. Natl. Cancer Inst.* 1994;86:1584-92.
7. Daling JR, Brinton LA, Voigt LF et al. Risk of breast cancer among white women following induced abortion. *Am. J. Epidemiol.* 1996;144(4):373-80.
8. Newcomb PA, Storer BE, Longnecker MP et al. Pregnancy termination in relation to risk of breast cancer. *J.A.M.A.* 1996 Jan 24-31;275(4):283-7.
9. Palmer JR, Rosenberg L, Rao RS et al, Induced and spontaneous abortion in relation to risk of breast cancer (United States). *Cancer Causes Control* 1997 Nov;8(6):841-9.
10. Nishiyama F. The epidemiology of breast cancer in Tokushima Prefecture. *Shikoku Ichi* 1982;38:333-43 (in Japanese).
11. Le M-G, Bachelot A, Doyon F, Kramar K, Hill C. Oral Contraceptive Use and Breast or Cervical Cancer: Preliminary Results of a French Case-Control Study. In: Wolff F-P, Scott JS, eds. *Hormones and Sexual Factors in Human Cancer Aetiology*. Amsterdam: Elsevier, 1984;139-47.
12. Lipworth L, Katsouyanni K, Ekborn A, Michels KB, Trichopoulos D. Abortion and the risk of breast cancer: A case-control study in Greece. *Int. J. Cancer* 1995;61:181-4.
13. Rookus MA, van Leeuwen FE. Induced abortion and risk for breast cancer: Reporting (recall) bias in a Dutch case-control study. *J. Natl. Cancer Inst.* 1996 Dec 4;88(23):1759-64.
14. Bu L, Voigt L, Yu Z, Malone K, Daling J. Risk of breast cancer associated with induced abortion in a population at low risk of breast cancer. *Am. J. Epidemiol.* 1995;141:S85 (abstract 337).
15. Talamini R, Franceschi S, La Vecchia C et al. The role of reproductive and menstrual factors in cancer of the breast before and after menopause. *Eur. J. Cancer* 1996;32A(2):303-10.
16. Luporsi E. Unpublished study (1988) data used in Andrieu N, Duffy SW, Rohan TE et al. Familial risk, abortion and their interactive effect on the risk of breast cancer: A combined analysis of six case-control studies. *Br. J. Cancer* 1995;72:744-51.
17. Rohan TE, McMichael AJ, Baghurst PA. A population-based case-control study of diet and breast cancer in Australia. *Am. J. Epidemiol.* 1988;128:478-89: data in Andrieu N, Duffy SW, Rohan TE et al. Familial risk, abortion and their interactive effect on the risk of breast cancer: A combined analysis of six case-control studies. *Br. J. Cancer* 1995;72:744-51.
18. Brind J, Chinchilli VM, Severs WB, Summy-Long J. Induced abortion as an independent risk factor for breast cancer: A comprehensive review and meta-analysis. *J. Epidemiol. Community Health* 1996;50:481-96.
19. Carroll PS. The breast cancer epidemic: Modeling and forecasts based on abortion and other risk factors. *Journal of American Physicians and Surgeons* 2007;12(3):72-8.
20. Ozmen V, Ozcinar B, Karanlik H et al. Breast cancer risk factors in Turkish women: A University Hospital based nested case control study. *World Journal of Surgical Oncology* 2009;7:37.

- doi:10.1186/1477-7819-7-37. Available from URL: <http://wjso.com/content/7/1/37> [Accessed 19 November 2009].
21. Xing P, Li J, Jin F. A case-control study of reproductive factors associated with subtypes of breast cancer in Northeast China. *Med. Oncol.* 2009 Sep 23. [Epub ahead of print]. Abstract available from URL: [http://www.ncbi.nlm.nih.gov/pubmed/19771534?itool=EntrezSystem2.PEntrez.Pubmed.Pubmed\\_ResuItsPanel.Pubmed\\_RVDocSum&ordinalpos=1](http://www.ncbi.nlm.nih.gov/pubmed/19771534?itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResuItsPanel.Pubmed_RVDocSum&ordinalpos=1) [Accessed 19 November 2009].
  22. Brind J. The abortion-breast cancer connection. *National Catholic Bioethics Quarterly* Summer 2005:303-29. Available from URL: [http://www.AbortionBreastCancer.com/Brind\\_NCBQ.PDF](http://www.AbortionBreastCancer.com/Brind_NCBQ.PDF) [Accessed 8 January 2009].
  23. Brind J. Induced abortion as an independent risk factor for breast cancer: A critical review of recent studies based on prospective data. *Journal of American Physicians and Surgeons* 2005;10(4):105-10. Available from URL: <http://www.jpands.org/vol10no4/brind.pdf> [Accessed 8 January 2009].
  24. Brind J. Induced abortion and breast cancer risk: A critical analysis of the report of the Harvard nurses study II. *Journal of American Physicians Surgeons* 2007;12(2):38-39.
  25. Brind J. Scientific Developments Relating to the Effect of Abortion on Risk of Future Breast Cancer: 2004 Oxford “reanalysis” of worldwide data. August 2007. Available from URL: <http://www.bcpinstitute.org/ABCUKParliamenttestimony-2007.htm> [Accessed 15 January 2010].
  26. Lanfranchi A. The abortion-breast cancer link revisited. *Ethics and Medics* November 2004;29(11):1-4.
  27. Furton E. The corruption of science by ideology. *Ethics and Medics* December 2004;29(12):1-2.
  28. Schlaflly AL. Legal implications of a link between abortion and breast cancer. *Journal of American Physicians and Surgeons* 2005;10(1):11-14. Available from URL: <http://www.jpands.org/vol10no1/aschlaflly.pdf> [Accessed 8 January 2009].
  29. DeLellis Henderson KD, Sullivan-Halley J, Reynolds P et al. Incomplete pregnancy is not associated with breast cancer risk: The California teachers study. *Contraception* 2008;77:391-6.
  30. Abortion Breast Cancer. Press Releases. California Study Denying Abortion-Cancer Link is Fraudulent, Says Scientist. Available from URL: [http://www.abortionbreastcancer.com/press\\_releases/080611](http://www.abortionbreastcancer.com/press_releases/080611) [Accessed 6 August 2008].
  31. Abortion Breast Cancer. Commentary on the Study DeLellis Henderson et al 2008. Available from URL: <http://www.abortionbreastcancer.com/news/080611> [Accessed 6 August 2008].
  32. Breast Cancer Prevention Institute. Reproductive Breast Cancer Risks and Breast Lobule Maturation. Online brochure. Available from URL: <http://www.bcpinstitute.org/reproductive.htm> [Accessed 17 January 2010].
  33. Breast Cancer Prevention Institute. The Biologic Cause of the Abortion Breast Cancer Link: The Physiology of the Breast. May, 2003. Fact sheet. Available from URL: <http://www.bcpinstitute.org/physiology.htm> [Accessed 17 January 2010].
  34. Breast Cancer Prevention Institute. Available from URL: <http://www.bcpinstitute.org/home.htm> [Accessed 15 January 2010].
  35. Houldsworth K. Ends of the Earth Productions. Abortionbreastcancer.com Commercial 1. YouTube. Available from URL: <http://www.youtube.com/watch?v=xj41i6dXPTY> [Accessed 15 January 2010].
  36. Dolle JM, Daling JR, White E et al. Risk factors for triple-negative breast cancer in women under the age of 45 years. *Cancer. Epidemiol. Biomarkers Prev.* 2009 Apr;18(4):1157-66. Epub 2009 Mar 31.